

ON SOME SIMPLE JACOBSON RADICAL RINGS

Ferenc SZÁSZ\*

Reprinted from MATHEMATICA JAPONICAE  
Vol. 18, No. 3, 1973

# ON SOME SIMPLE JACOBSON RADICAL RINGS

Ferenc SZÁSZ\*

(Received January 7, 1972)

It is well known, that among the various concrete radicals of associative rings the Jacobson radical seems to be the most useful for investigation of the structure of rings (see N. Jacobson [2]). On the other side, for much structure theorems for rings, using subdirect sums and some concrete radicals, the so-called  $F$ -radicals by B. Brown and N. H. McCoy [1], the subdirectly irreducible rings play an important rôle. Among these rings the simple rings (i. e. the rings  $A$  with  $A^2=A$  and without nontrivial ideals) are of extraordinary significance.

It is well known (cf. author [7]), that every simple Jacobson radical ring with minimum condition on principal right ideals is the ring  $(O)$ , since it satisfies, by author [7] necessarily  $A^2=(O)$ , whence  $A^2=A$  implies  $A=(O)$ , indeed. E. Sasiada [6] has constructed a nontrivial simple Jacobson radical ring. Jacobson's famous problem asks:

Does there exist nontrivial simple nil rings?

This problem is yet unsolved, but some interesting properties of these rings were asserted by W. A. McWorter [5]. For some characterizations of the Jacobson radical we refer yet e. g. to Chapter 5 of the book of A. Kertész [3], Chapter 4 of the book of author [9] and author [8]. For some properties of simple Jacobson radical rings we refer to author [10].

According to A. G. Kurosh's book [4], every group  $G$ , written multiplicatively, can be embedded into a group  $\bar{G}$  such that every two elements  $x, y \in \bar{G}$ , which are different from  $1 \in \bar{G}$ , also are conjugate in  $\bar{G}$ , that is there exists an element  $z \in \bar{G}$  satisfying  $x = z^{-1}yz$ .

Considering this grouptheoretical property, we understand by an  $\Omega$ -ring  $A$  a Jacobson radical ring, all nonzero elements of which are quasiconjugate, i. e. if  $x \neq 0, y \neq 0$ , and  $x, y \in A$ , then there exists an element  $z \in A$  such that

$$(*) \quad x = (1-z)y(1-z)^{-1}$$

**Proposition 1.** *Every  $\Omega$ -ring satisfying  $A^2=A$ , is a simple Jacobson radical ring.*

*Proof.* It is sufficient, by the assumed Jacobson radical property, only shown the simplicity of  $A$ , which follows from  $A^2=A$  and from  $x \in (y)$ , the principal

---

\* Mathematical Institute of Hungarian Academy of Sciences, Budapest, Hungary.

ideal generated by  $y$  in  $A$ , since  $(1-z)^{-1}$  has obviously a form  $1-w$  with  $w \in A$ , and  $z+w-zw=0$ .

**Proposition 2.** *Every  $\Omega$ -ring, which satisfies  $A^2=A$ , is an algebra over a prime field  $F_p$  ( $p=0$  or a prime number).*

*Proof* is trivial, since either  $pA=0$  for a suitable prime number  $p$ , or  $pA=A$  holds for every  $p$ . In latter case  $A$  is of characteristic 0.

**Proposition 3.** *Every  $\Omega$ -ring, which contains nonzero nilpotent elements, satisfies  $A^2=0$  and it consists only of two elements:  $A=[0, a]$  with  $a^2=a+a=0$ .*

*Proof.* If  $A$  has nonzero nilpotent elements, then there exists an  $a \neq 0$  with  $a^2=0$ , whence by (\*) every element of  $A$  has a square zero, and thus  $bc=-cb$  holds for every  $b, c \in A$ . Furthermore  $(1-z)^{-1}=1+z$ , whence (\*) and  $zy=-yz$ ,  $z^2=0$  imply

$$x=(1-z)y(1+z)=y+2yc \in (y)_r,$$

the principal right ideal generated by  $y$  in  $A$ . Thus  $A$  coincides with the minimal right ideal  $(y)_r$ , which satisfies  $A^2=0$ . This implies  $|A|=2$ , indeed.

**Proposition 4.** *Every  $\Omega$ -ring without nonzero nilpotent elements does not contain nonzero divisors of zero.*

*Proof.* Assume  $xy=0$  and  $x \neq 0$ . Then, by our assumption,  $(yAx)^2=0$  and  $(yx)^2=0$  we have  $yAx=yx=0$  and thus also  $xAy=0$ . This yields:

$$(x).(y)=(y).(x)=0,$$

which by the definition at (\*) of  $\Omega$ -rings implies in case  $y \neq 0$  by  $(x)=(y)$  the contradiction  $x^2=0$ ,  $x=0$ .

**Theorem 5.** *If  $A$  is an  $\Omega$ -ring without nonzero divisors of zero, then every its element is transcendental over the prime field  $F_p$  belonging to  $A$  (according to Proposition 2). Furthermore, for every polynomial  $f(x) \in x, \mathfrak{I}[x]$  (here  $\mathfrak{I}$  denotes the ring of rational integers) and for every nonzero element  $a \in A$  there exists an element  $b$ , which satisfies  $a=f(b)$  ( $b \in A$ )*

*Proof.* By Chapter I of N. Jacobson [2] the elements of the Jacobson radical of an algebra over a field are either nilpotent or transcendental over  $F_p$ , but our assumption excludes the nilpotent elements. Thus  $A$  is absolutely transcendental over  $F_p$ . Therefore, for every polynomial  $f(x) \in x, \mathfrak{I}[x]$  and every nonzero element  $a \in A$  we have  $f(a) \neq 0$ . According to the definition at (\*) there exists an element  $c \in A$  such that  $f(a)=(1-c)a(1-c)^{-1}$ . Let us define the desired element  $b \in A$  by  $b=(1-c)^{-1}a(1-c)$ . Then

$$\varphi: x \rightarrow (1-c)^{-1}x(1-c)$$



where  $x \in A$ , is an automorphism of the ring, whence

$$f(b) = f(\varphi a) = \varphi(f(a)) = (1-c)^{-1}f(a)(1-c) = a$$

holds, indeed.

**Corollary 6.** ("Fermat's strong theorem", for  $\Omega$ -rings). *Let  $A$  be an  $\Omega$ -ring without nonzero nilpotent elements, furthermore let  $n \geq 3$  be a natural number and  $a$  and  $b$  be arbitrary elements of  $A$  such that  $a^n + b^n \neq 0$ . Then there exists an element  $c \in A$  such that  $a^n + b^n = c^n$  holds.*

*Proof.* By our assumption, Proposition 4 and Theorem 5, the algebra  $A$  over  $F_p$  is absolutely transcendental. Put  $f(x) = x^n \in x.\mathfrak{F}[x]$ , where  $n \geq 3$ . If  $a^n + b^n \neq 0$ , then an application of Theorem 5 to this case completes the proof.

**Remark 7.** The assertion of Corollary 6 and its proof are true naturally also for  $n=2$ , and the case  $n=1$  is trivial, but these cases naturally differ from Fermat's conjecture for the ring  $\mathfrak{F}$ .

**Proposition 8.** *In every  $\Omega$ -ring  $A$  without nonzero nilpotent elements the centralizer  $C_x$  of an arbitrary nonzero element  $x \in A$  is properly larger than the subring  $\{x, x'\}$  generated by  $x$  and  $x'$ , where  $x' \in A$  and  $x + x' - x.x' = 0$ .*

*Proof.* We have  $(1-x)(1-x') = 1$ . Furthermore, there exists an element  $z \in A$  with  $x' = (1-z)x(1-z)^{-1}$ . Therefore  $(1-x')(1-(1-z)x'(1-z)^{-1}) = 1$ , whence, by the unicity of quasiinverses, follows

$$x = (1-z)x'(1-z)^{-1},$$

consequently  $x = (1-z)^2 x (1-z)^{-2}$  and  $x' = (1-z)^2 x' (1-z)^{-2}$ . In other words,  $c = z^2 - 2z$  belongs to the centralizers of the subrings  $\{x\}$  and  $\{x'\}$ , but, by the absolute transcendence of  $A$  over  $F_p$  holds  $x \notin \{x'\}$  and  $x' \notin \{x\}$ , furthermore  $c = z^2 - 2z \notin \{x, x'\}$ . Therefore, the centralizer  $C_x$  of  $x$  is properly larger, than  $\{x, x'\}$ , indeed, since  $C_x \supseteq \{x, x'\}$

*Problem.* Does there exist an absolutely transcendental  $\Omega$ -ring?

#### References

- [1] B. Brown-N. H. McCoy, Radicals and subdirect sums, Amer. Jour. Math. **69**, 46-58 (1947).
- [2] N. Jacobson, Structure of Rings (2. edition) Providence (1964).
- [3] A. Kertész, Vorlesungen über Artinsche Ringe, Budapest, Akadémiai Kiadó (1968)
- [4] A. G. Kurosh, Theory of Groups. (3. edition), Moscow (1967, in Russian)
- [5] W. A. McWorter, Some simple properties of simple nil rings, Canadian Math. Bulletin **9**: 2, 197-200 (1966).
- [6] E. Sasiada, Solution of the problem of existence of simple radical ring, Bull. Acad. Polon. Sci. **9**, 257 (1961)
- [7] F. Szász, Ueber Ringe mit Minimalbedingung für Hauptrechtsideale, I, Publ. Math. Debrecen **7**, 54-64 (1960); II, Acta Math. Acad. Sci. Hungar. **12**, 417-439 (1961); III, Acta Math. Acad. Sci. Hungar. **14**, 447-461 (1963).

- [ 8 ] F. Szász, Further characterizations of the Jacobson radical, Proc. Japan Acad. (to appear)
- [ 9 ] F. Szász, Radikale der Ringe, Budapest, Akadémiai Kiadó. (to appear)
- [10] F. Szász, Beiträge zur Radikaltheorie der Ringe, Publicationes Math. Debrecen 17 : 1-4, 267-272 (1970).